

Rye

Publications

2008

Heiniö, R.-L., Myllymäki O., Pihlava, J.-M, Adlercreutz, H., Heinonen, S.-M., Liukkonen K.-H. and Poutanen, K. Quantities of phenolic compounds and their impacts on the perceived flavour attributes of rye grain. *J Cereal Sci*, in press.

Kallio, P., Kolehmainen, M., Laaksonen, D.E., Pulkkinen, L., Atalay, M., Mykkänen, H., Uusitupa, M., Poutanen, K., Niskanen, L. Inflammation markers are modulated by responses to diets differing in postprandial insulin responses in individuals with the metabolic syndrome. *Am J Clin Nutr* 87, 1497-1503.

Kamal-Eldin, A., Åman P., Zhang, J.-X., Bach Knudsen, K.-E., Poutanen, K. High-fibre products: rye bread and other rye products. In: *Technology of Functional Cereal Products*. Hamaker, B.R. (Ed.), Woodhead Publishing Ltd, Abington, Cambridge, England, pp. 233–260.

Seal, C.J., de Mul, A., Eisenbrand, G., Haverkort, A.J., Franke, K., Lalljie, S.P.D., Mykkänen H., Reimerdes, E., Scholz, G., Somoza, V., Tuijtelars, S., van Boekel, M., van Klaveren, J., Wilcockson, S.J., Wilms, L. Risk-benefit considerations of mitigation measures on acrylamide content of foods - a case study on potatoes, cereals and coffee. *Br J Nutr* 99, S1-S46.

2007

Aura, A., Myllymäki, O., Bailey, M., Penalvo, J.-L., Adlercreutz, H. and Poutanen, K. Interrelationships between carbohydrate type, phenolic acids and initial pH on in vitro conversion of enterolactone from rye lignans. *Dietary Fibre Components and Functions*. Salovaara, H.; Gates, F.; Tenkanen, M. (Eds.), Wageningen Academic Publishers, The Netherlands, pp. 235–245.

Autio, K., Liukkonen, K.-H., Mykkänen, H., Katina, K., Roininen, K. and Poutanen, K. Structure of whole grain breads: Sensory perception and health effects. In: *Whole Grains and Health* (Marquart L., Jacobs D.R., McIntosh G.H., Poutanen K., Reicks M., eds.), Blackwell Publishing, p. 115-122.

Gråsten, S.M., Juntunen, K.S., Mättö, J., Mykkänen, O.T., El-Nezami, H., Adlercreutz, H.C.T., Poutanen, K.S. and Mykkänen, H.M. High-fiber rye bread improves bowel function but does not cause other putatively positive changes in the metabolic activity of intestinal microbiota. *Nutr Res* 27, 454–461.

Kallio, P., Kolehmainen, M., Laaksonen, D.E., Kekäläinen, J., Salopuro, T., Sivenius, K., Pulkkinen, L., Mykkänen, H., Niskanen, L.K., Uusitupa, M.I. and Poutanen, K. Dietary carbohydrate modification induces marked alterations in gene expression of white adipose tissue in persons with metabolic syndrome – the FUNGENUT Study. *Am J Clin Nutr* 85, 1417–1427.

Katina, K., Liukkonen, K.-H., Kaukovirta-Norja, A., Adlercreutz, H., Heinonen, S.-M., Lampi, A.-M., Pihlava, J. and Poutanen, K. Fermentation induced changes on the nutritional value of native or germinated rye. *J Cereal Sci* 46, 348–355.

Katina, K., Laitila, A., Juvonen, R., Liukkonen, K.-H., Kariluoto, S., Piironen, V., Landberg, R., Åman, P and Poutanen, K. Bran fermentation as a means to enhance technological properties and bioactivity of rye. *Int J Food Microbiol* 24,175–186.

Nyström, L., Lampi, A.-M., Rita, H., Aura, A.-M., Oksman-Caldentey, K.-M. and Piironen, V. Effects of processing on availability of total plant sterols, steryl ferulates and steryl glycosides from wheat and rye bran. *J Agric Food Chem* 24, 9059–9065.

2006

Hallikainen, M., Toppinen, L., Mykkänen, H., Ågren, J., Laaksonen, D., Niskanen, L., Miettinen, T.A., Poutanen, K. and Gylling, H. Interaction between cholesterol and glucose metabolism during dietary carbohydrate modification in subjects with the metabolic syndrome. *Am J Clin Nutr* 84, 1385–1392.

Hätönen, K.A., Similä, M.E., Virtamo, J.R., Eriksson, J.G., Hannila, M.-L., Sinkko, H.K., Sundvall, J.E., Mykkänen, H.M. and Valsta, L.M. Methodologic considerations in the measurement of glycemic index: glycemic response to rye bread, oatmeal porridge, and mashed potato. *Am J Clin Nutr* 84, 1055–1061.

Liukkonen, K.-H., Heiniö, R.-L., Salmenkallio-Marttila, M., Autio, K., Katina, K. and Poutanen, K. Rye. In: *Bakery products: Science and Technology*, Hui, Y.H. et al. (Eds). Blackwell Publishing. Ames, Iowa, USA, pp. 109–122.

Liukkonen, K.-H., Katina, K., Kaukovirta-Norja, A., Lampi, A.-M., Kariluoto, S., Piironen, V., Heinonen, S.-M., Adlercreutz, H., Nurmi, A., Pihlava, J.-M. and Poutanen, K. Influence of germination conditions on the bioactivity of rye. In: *Whole Grains and Health*. L. Marquardt, M. Reicks, D. Jacobs, G.H. McIntosh, K. Poutanen (Eds.), Blackwell Publishing, Oxford, pp. 229–239.

Poutanen, K., Laaksonen, D., Autio, K., Mykkänen, H. and Niskanen, L. The role of carbohydrates in the prevention and management of type 2 diabetes. In: *Functional Food Carbohydrates* (Biliaderis, C., Isidczyk, M., Eds), CRC Press, p. 387-411.

2005

Aura A.-M., Karppinen, S., Virtanen, H., Forssell, P., Heinonen, S., Nurmi, T., Adlercreutz, H. and Poutanen, K. Processing of rye bran influences both the fermentation of dietary fibre and the bioconversion of lignans by human faecal flora in vitro. *J Sci Food Agr* 85, 2085–2093.

Laaksonen, D.E., Toppinen, L.K., Juntunen, K.S., Autio, K., Poutanen, K.S., Niskanen, L.K. and Mykkänen, H.M. Dietary carbohydrate modification enhances insulin secretion in individuals with the metabolic syndrome. *Am J Clin Nutr* 82, 1218–1227.

Linko, A., Juntunen, K., Mykkänen, H. and Adlercreutz, H. Whole-grain rye bread consumption by women correlates with plasma alkylresorcinols and increases their concentration compared with low-fiber wheat bread. *J Nutr* 135, 580–583.

2004

Kariluoto, S., Vahteristo, L., Salovaara, H., Katina, K., Liukkonen, K.-H. and Piironen, V. Effect of baking method and fermentation on folate content of rye and wheat breads. *Cereal Chem* 81, 134–139.

2003

Heiniö, R.-L., Katina, K., Wilhelmson, A., Myllymäki, O., Rajamäki, T., Latva-Kala, K., Liukkonen, K.-H. and Poutanen, K. Relationship between sensory perception and flavour-active volatile compounds of germinated, sourdough fermented and native rye following the extrusion process. *Food Sci Technol* 36, 533–545.

Heiniö, R-L., Liukkonen, K.-H., Katina, K., Myllymäki, O. and Poutanen, K. Milling fractionation of rye produces different sensory profiles of both flour and bread. *Food Sci Technol* 36, 577–583.

Juntunen, K.S., Laaksonen D.E., Poutanen K.S., Niskanen, L.K., and Mykkänen H.M. High-fiber rye bread and insulin secretion and sensitivity in healthy postmenopausal women. *Am J Clin Nutr* 77, 385–391.

Juntunen, K., Laaksonen, D.E., Autio, K., Niskanen, L.K., Holst, J.J., Savolainen, K.E., Liukkonen, K.-H., Poutanen, K. and Mykkänen, H. Structural differences between rye and wheat bread but not total fiber content may explain the lower postprandial insulin response to rye bread. *Am J Clin Nutr* 78, 957–964.

Karppinen, S., Myllymäki, O., Forssell, P. and Poutanen K. Fructan content of rye and rye products. *Cereal Chem* 80, 168–171.

Liukkonen, K.-H., Katina, K., Wilhelmsson, A., Myllymäki, O., Lampi, A., Kariluoto, S., Piironen, V., Heinonen, S., Nurmi, T., Adlercreutz, H., Peltoketo, A., Pihlava, J., Hietaniemi, V. and Poutanen, K. Session: Health effects of whole grains. Process-induced changes on bioactive compounds in whole grain rye. *Proc Nutr Soc* 62, 117–122.

Oikarinen, S., Heinonen, S., Karppinen, S., Mättö, J., Adlercreutz, H., Poutanen, K. and Mutanen, M. Plasma enterolactone or intestinal Bifidobacterium levels do not explain adenoma formation in multiple intestinal neoplasia (min) mice fed with two different types of rye bran fractions. *Br J Nutr* 90, 119–125.

2002

Gråsten, S.M., Pajari, A.-M., Liukkonen, K.-H., Karppinen, S. and Mykkänen, H.M. Fibers with different solubility characteristics alter similarly the metabolic activity of intestinal microbiota in rats fed cereal brans and inulin. *Nutr Res* 22:1435-1444.

Juntunen, K.S., Niskanen, L.K., Liukkonen, K.H., Poutanen, K.S., Holst, J.J. and Mykkänen, H.M. Postprandial glucose, insulin, and incretin responses to grain products in healthy subjects. *Am J Clin Nutr* 75, 254–262.

2001

Autio, K., Simoinen, T., Suortti, T., Salmenkallio-Marttila, M., Lassila, K. and Wilhelmson, A. Structural and enzymic changes in germinated barley and rye. *J Inst Brew* 107, 19–25.

Heinonen, S., Nurmi T., Adlercreutz H., Wähälä K., Liukkonen K., Poutanen K., Deyama T. and Nishibe S. In vitro metabolism of plant lignans: New precursors of mammalian lignans enterolactone and enterodiol. *J Agric Food Chem* 49, 3178–3186.

Karppinen, S., Kiiliäinen, K., Liukkonen, K., Forssell, P. and Poutanen, K. Extraction and in vitro fermentation of rye bran fractions. *J Cereal Sci* 34, 269–278.

Prättälä, R., Helasoja, V. and Mykkänen, H.M. The consumption of rye bread and white bread as dimensions of health life style in Finland. *Public Health Nutr* 4: 813-818.

Salovaara, H. and Autio, K. Rye and Triticale. In: *Cereals and Cereal Products, Chemistry and Technology*. Dendy, S.A.V. and Dobraszczyk, B.J. (Eds.), Aspen Publishers, Gaithersburg, Maryland, pp. 391–410.

2000

Gråsten, S., Leinonen, K., Poutanen, K., Gylling, H., Miettinen, T. and Mykkänen, H. Rye bread improves bowel function and decreases the concentration of some putative colon cancer risk markers in middle-aged women and men. *J Nutr* 130, 2215–2221.

Juntunen, K.S., Mazur, W.M., Liukkonen, K.-H., Poutanen, K.S., Adlercreutz, C.H.T. and Mykkänen, H.M. Consumption of wholemeal rye bread increases serum concentrations and urinary excretion of enterolactone compared with consumption of white wheat bread in healthy Finnish men and women. *Br J Nutr* 84, 839–846.

Karppinen, S., Liukkonen, K., Aura, A.-M., Forssell, P. and Poutanen, K. In vitro fermentation of polysaccharides of rye, wheat and oat brans and inulin by human faecal bacteria. *J Sci Food Agric* 80, 1469–1476.

Leinonen, K.S., Poutanen, K. and Mykkänen, H.M. Rye bread decreases serum total and LDL cholesterol in men with moderately elevated serum cholesterol. *J Nutr* 130, 164–170.

Nilsson, M., Andersson, R., Andersson, R. E., Autio, K. and Åman, P. Heterogeneity in a water-extractable rye arabinoxylan with a low degree of disubstitution. *Carboh Polym* 41, 397–405.

Ouwehand, A.C., Gråsten, S., Niemi, P., Mykkänen, H. and Salminen, S. Wheat or rye supplemented diets do not affect faecal mucus concentrations or the adhesion of probiotic microorganisms to faecal mucus. *Lett Appl Microbiol* 31:30-33.

Poutanen, K. and Liukkonen, K. Rye: a Nordic focus. *Functional Foods* 3, 30–32.

Turpeinen, A., Juntunen, K., Mutanen, M. and Mykkänen, H. Wholemeal rye bread does not differ from low-fiber wheat bread on the effects on hemostatic factors. *Eur J Clin Nutr* 54:418-423.

1999

Autio, K., Flander, L., Heinonen, R. and Kinnunen, A. Comparison of small and large deformation measurements of whole meal rye doughs. *Cereal Chem* 76, 912–914.

Aura, A-M, Härkönen, H., Fabritius, M. and Poutanen, K. Development of an enzymatic in vitro method and assessment of its performance using wheat and rye breads. *J Cereal Sci* 29, 139–152.

Leinonen, K., Liukkonen, K., Poutanen, K., Uusitupa, M. and Mykkänen, H. Rye bread decreases postprandial insulin response but does not alter glucose response in healthy Finnish subjects. *Eur J Clin Nutr* 53, 262–267.

Liukkonen, K., Poutanen, K., Leinonen, K. and Mykkänen, H. Ruistuotteet suosiossa (Rye products gain popularity) *Sairaala-lehti* 61, 26–28.

Poutanen, K. and Liukkonen, K-H. Focus on Rye. *Ingredients, Health & Nutrition* 2, 14–16.

Poutanen, K. Functional Rye Products. *Fi Food Summit Proceedings 1999*, 14–16 September, Paris 1999.

1998

Autio, K., Fabritius, M. and Kinnunen, A. Effect of germination and water content on the microstructure and rheological properties of two rye doughs. *Cereal Chem* 75, 10–14.

Poutanen, K., Liukkonen, K., Härkönen, H., Mazur, W. and Adlercreutz, H. Miksi ruisleipä on terveellistä (Chemistry behind the health benefits of rye), *Kemia-Kemi* 25, 235–238.

1997

Autio, K., Parkkonen, T. and Fabritius, M. Observing structural differences in wheat and rye breads. *Cereal Foods World* 8, 702–705.

Fabritius, M., Gates, F., Salovaara, H. and Autio, K. Structural changes in insoluble cell walls in wholemeal rye doughs. *Lebensm-Wiss u Technol* 30, 367–372.

Hallmans, G., Zhang, J-X., Lundin, E., Landström, M., Åman, P., Adlercreutz, H., Härkönen, H. and Bach Knudsen, K. E. Influence of rye bran on the formation of bile acids and bioavailability of lignans. *Cereal Foods World* 8, 696–701.

Heiniö, R-L., Urala, N., Vainionpää, J., Poutanen, K. and Tuorila, H. Identity and overall acceptance of two types of sour rye bread. *Int J Food Sci Technol* 32, 169–178.

Härkönen, H., Pessa, E., Suortti, T. and Poutanen, K. Distribution and some properties of cell wall polysaccharides in rye milling fractions. *J Cereal Sci* 26, 95–104.

Nilsson, M., Åman, P., Härkönen, H., Hallmans, G., Bach Knudsen, K-E., Mazur, W. and Adlercreutz, H. Content of Nutrients and Lignans in Roller Milled Fractions of Rye. *J Sci Food Agric* 73, 143–148.

Nilsson, M., Åman, P., Härkönen, H., Mazur, W. and Adlercreutz, H. Nutrient and lignan content, dough properties and baking performance of Scandinavian rye samples. *Acta Agriculturae Scandinavica Section BB. Plant Science* 47, 26–34.

Parkkonen, T., Heinonen, R. and Autio, K. A new method for determining the area of cell walls in rye doughs based on fluorescence microscopy and computer-assisted image analysis. *Lebensm-Wisst u Technol* 30, 743–747.

Poutanen, K. Rye Bread: Added value in the world's bread basket. *Cereal Foods World* 8, 682–683.

1996

Autio, K., Härkönen, H., Parkkonen, T., Frigård, T., Siika-aho, M., Poutanen, K. and Åman, P. Effects of purified endo- β - xylanase and endo- β - glucanase on the structural and baking characteristics of rye doughs. *Lebensm-Wiss u technol* 29, 18–27.

Juokslahti, T., Parkkonen, T. and Bedford, M. The effect of enzyme supplementation of chicken rye feed on the viscosity and microstructure of digesta. *Zootechnika* XLI 297, 89–98.

Pettersson, D, Åman, P., Bach Knudsen, K-E, Lundin, E., Zhang, J-X., Hallmans, G., Härkönen, H. and Adlercreutz, H. Intake of Rye Bread by Ileostomists Increases Ileal Excretion of Fiber Polysaccharide Components and Organic Acids but Does Not Increase not Plasma or Urine Lignans and Isoflavonoids. *J Nutr* 126, 1594–1600.

Salminen, A., Latva-Kala, K., Randell, K., Hurme, E., Linko, P. and Ahvenainen, R. The effect of ethanol and oxygen absorption on the shelf-life of packed sliced rye bread. *Packaging Tech Sci* 9, 29–42.

1995

Autio, K., Koskinen, M. and Parkkonen, T. Structural changes in cell walls of rye doughs during incubation. International rye symposium: Technology and products, Helsinki, 7-8 Dec. 1995. Poutanen, K. and Autio, K. (eds.), Espoo 1995. VTT Symposium 161, pp. 141–154.

Parkkonen, T., Härkönen, H. and Autio, K. Effect of baking on the microstructure of rye cell walls and protein. Cereal Chem 1, 58–63.

Poutanen, K. and Autio, K. (Eds.) International rye symposium: Technology and products, Helsinki, 7-8 Dec. 1995. Espoo 1995. VTT Symposium 161, 221 p.

Poutanen, K., Härkönen, H., Laurikainen, T., Suortti, T., Autio, K., Luonteri, E. and Tenkanen, M. Enzymatic accessibility of rye cell wall polysaccharides. International rye symposium: Technology and products, Helsinki, 7-8 Dec. 1995. Poutanen, K. and Autio, K. (Eds.), Espoo 1995. VTT Symposium 161, 121–130.

1994

Härkönen, H., Lehtinen, P., Suortti, T., Parkkonen, T., Siika-Aho, M. and Poutanen, K. The effects of a xylanase and a β -glucanase from *Trichoderma reesei* on the non-starch polysaccharides of whole meal rye slurry. J Cereal Sci 21, 173–183.

1993

Härkönen, H., Autio, K., Parkkonen, T., Åman, P. and Poutanen, K. The role of pentosans in rye bread baking. Proceedings from the 25th Nordic Cereal Congress: the Nordic Cereal Industry in an Integrating Europe, Helsinki, 6-9 June 1993.

Doctoral theses

Heiniö, R.-L. Influence of processing on the flavour formation of oat and rye. VTT, 2003.
Pdf: <http://www.vtt.fi/inf/pdf/publications/2003/P494.pdf>

Juntunen, K. Rye bread and glucose and insulin in healthy adults. University of Kuopio, 2003.
Pdf: <http://www.uku.fi/vaitokset/2003/isbn951-781-353-8.pdf>

Karppinen S. Dietary fibre components of rye bran and their fermentation in vitro. VTT, 2003.
Pdf: <http://www.vtt.fi/inf/pdf/publications/2003/P500.pdf>